FLOWERING SEASONS OF AUSTRALIAN PLANTS.

No. II.

BY E. HAVILAND, F.L.S.

PLANTS FLOWERING IN THE NEIGHBOURHOOD OF SYDNEY DURING THE MONTH OF AUGUST, IN ADDITION TO THOSE ENUMERATED IN THE LIST FOR JULY, NEARLY ALL OF WHICH ARE STILL IN FLOWER.

As it is impossible for the author to visit more than a few localities during each month, these lists must necessarily, for the present, be very imperfect. Supplementary lists will, however, be added as each month recurs, until, it is hoped, the flowering seasons of at least the Cumberland plants have been ascertained. When sufficient data have been collected, a tabulated statement will be published, giving against each species its complete flowering season.

Ranunculaceæ—

Clematis aristata.

Dilleniaceæ—

Hibbertia volubilis

dentata.

Polygaleæ—

Comesperma ericinum.

Tremandreæ—

Tetratheca pilosa.

Rutaceæ—

Phebalium denticulatum.

Euphorbiaceæ—

Amperea spartioides Poranthera microphylla. Leguminosæ—

Gompholobium Huegelii Pultenea echinulata Acacia elongata Bossiwa ensata Indigofera australis Oxylobium trilobatum.

Crassulaceæ—

Tillea purpurata.

Myrtaceæ—

Callistemon rigidus ,, linearis Melaleuca ericifolia.

Umbelliferæ—	Epacrideæ—
Trachymene incisa	Epacris pulchella
Proteaceæ—	,, purpurascens
Hakea acicularis	Brachyloma daphnoides
$Conospermum\ longifolium$	Leucopogon amplexicaulis
var. angustifoliu	m. ,, esquamatus
Compositæ—	,, lanceolatus
$Gnaphalium\ Japonicum$,, biflorus.
$Epaltes\ australis$	Orchideæ—
Senecio australis.	Diuris sulphurea
Campanulaceæ—	Dendrobium speciosum
Lobelia gracilis	,, linguiforme
,, anceps	$Caladenia\ cœrulea$
,, dentata	,, carnea
Wahlenbergia gracilis.	Cyrtostylis reniformis
Goodeniaceæ—	Lyperanthus suaveolens
Dampiera Brownii.	,, nigricans
Myrsinaceæ—	Pterostylis pedunculata
Myrsine variabilis.	,, concinna
Scrophularineæ—	,, $curta$
Euphrasia speciosa.	,, reflexa.
Bignoniaceæ—	Liliaceæ—
$Tecoma\ australis.$	$Stypandra\ umbellata$

Stypandra umbellata ,, glauca.